arXiv:1304.1744v1 [physics.acc-ph] 5 Apr 2013

Fundamental Photoemission Brightness Limit from Disorder Induced Heating

Jared Maxson* and Ivan Bazarov
Cornell Laboratory for Accelerator-Based Sciences and Education,
Cornell University, Ithaca, New York 14853

Christopher Coleman-Smith
Department of Physics, Duke University, Durham, NC 27708

Weishi Wan and Howard Padmore
Advanced Light Source, Lawrence Berkeley National Lab,
1 Cyclotron Road MS 2R0400, Berkeley, CA 94720

(Dated: April 8, 2013)

Abstract

We determine the limit of the highest achievable photoemitted electron brightness, due to heating just after emission into vacuum, applicable to both relativistic and nonrelativistic photoelectron production. This heating is due to poorly screened Coulomb interactions equivalent to disorder induced heating seen in ultracold neutral plasmas. Owing to the multi-scale nature of the initial interactions of such systems, an N-body tree algorithm is used to compute the universal scaling of the disorder induced heating in fully contained bunches, and is shown to agree well with a simple model utilizing the tabulated correlated energy of one component plasmas. We also present simulations for beams undergoing Coulomb explosion at the photocathode, and demonstrate that both the temperature growth and subsequent cooling must be characterized by correlated effects, as well as correlation-frozen dynamics. In either case, the induced temperature is found to be of several meV for typical photoinjector beam densities, a significant fraction of the intrinsic beam temperature of the coldest semiconductor photocathodes.

Great progress has been made in the cooling of photoemitted electrons, in particular by those working with negative electron affinity semiconductor photocathodes and groups pursuing electron emission from laser cooled atoms. Several semiconductor photocathode groups have measured photoemission of equivalent temperatures well below $kT = 25$ meV, or the thermal energy at room temperature, either via the cryogenic cooling of the photocathode [1], or via the maintenance of a pristine photoemissive conditions in ultra high vacuum, allowing the low effective mass of conduction electrons to produce an effect equivalent to Snell's law in which the velocity spread of electrons is drastically reduced at the vacuum interface [2]. Furthermore, using laser cooled atoms which are photoionized, there has been production of electrons at temperatures equivalent to ~ 1 meV [3].

This temperature of emission, along with the accelerating field at the source E_{acc}, are the only two parameters of the fundamental brightness (transverse, normalized) limit of photoinjectors [4], such that:

$$\mathcal{B}_{\mathrm{n,4D}}\bigg|_{\mathrm{max}} = \frac{mc^2 f \epsilon_0 E_{\mathrm{acc}}}{2\pi kT} \tag{1}$$

where f is the bunch repetition rate. many next-generation photoinjector and ultrafast electron diffraction laboratories, whose applications depend critically on beam brightness, plan to merge such ultra cold photoemission sources with photoguns to produce relativistic particle bunches. For various applications, such bunches may contain from $10^6 \rightarrow 10^9$ electrons, of densities in the range of $10^{17} \rightarrow 10^{20}$ m^{-3}.

The vast majority of all particle accelerator beam dynamics simulations and analytic calculations for relativistic photoinjectors rely on the space charge (SC) approximation to calculate the interparticle interaction, including nonrelativistic dynamics at the photocathode. Within the SC approximation, the Debye length of the electrons is assumed to be much larger than the average interparticle separation. Analytically, this allows the reduction of the full 6N dimensional particle distribution function $\rho_{6N}(\mathbf{r}_1, \mathbf{v}_1, \mathbf{r}_2, ..., \mathbf{r}_n, \mathbf{v}_n)$ to a single particle form $\rho_6(\mathbf{r}, \mathbf{v})$. In beam dynamics simulations, this allows for the use of macroparticles representing many electrons, from which the self-consistent calculation and evolution of the mean field corresponding macroparticle distribution is made.

However, for the lowest temperatures attained in semiconductor photoemission $kT \sim 5$ meV [2], and the lowest of the above densities, the Debye length is already on the order of the interparticle separation. Thus the space charge approximation is no longer valid. For larger densities, we expect the interaction lengths of the interparticle force to be much greater than the Debye length. Thus we expect some intrinsic velocity spread from the stochastic, poorly screened Coulomb forces. Further reduction of the photocathode emission temerature at sufficiently high density would be blurred out by this stochastic heating. In this Letter, we calculate the amount of this stochastic heating.

Though certain effects of Coulomb collisions are well studied in beam physics, as in the Boersch effect and intrabeam scattering, this high density, low temperature limit represents a new regime for beam calculations. Electron collisions in beams are traditionally handled via the diffusive Fokker-Planck method [5] or via averaging over two particle interactions [6], which both involve the reduction of the phase space dimensionality, as in the SC approximation. Thus, such methods are inadequate to describe the fully-coupled initial conditions of a near-zero temperature bunch.

The heating associated with the relaxation of a random, near-zero temperature distribution of charges is known to the ultracold neutral plasma (UNP) community. In such systems, a cold gas is laser ionized, and after a time on the order of the $\tau = 2\pi\omega_p^{-1} = 2\pi\left(n_0 e^2/m_e\epsilon_0\right)^{-1/2}$, the ion temperature is seen to rise significantly due to near-neighbor interactions, to nearly $3kT_c/2 = e^2/4\pi\epsilon_0 a$, where a is the Wigner-Seitz radius, $a = (3/4\pi n_0)^{1/3}$. In the traditional plasma physics terminology, this effect is referred to as disorder induced heating (DIH), and the effect is seen for plasma coupling parameters $\Gamma = e^2/4\pi\epsilon_0 akT$ of order unity or larger [7]. If we here assume a photocathode emission temperature of 5 meV, the above beam densities correspond to $\Gamma = 0.2 \rightarrow 2$. This estimate suggests that for the next generation of colder and denser electron beams, DIH serves as a hard limit to the achievable beam brightness.

Disorder induced heating has been experimentally observed for the ions in a neutral plasma [8]. In cold neutral plasmas, the electrons equilibrate much faster than the ionic ω_p^{-1}, and then serve to screen ion-ion interaction. An expression for the amount of DIH in ions was first given for Yukawa systems (for which electron screening, but not electron-ion recombination, is considered) in [9]. The initial ionic distribution is uncorrelated, and can be defined as the zero energy state. As the ions relax, the ion distribution begins to develop order, and the resulting correlations have an associated binding energy. This correlation binding energy can be calculated from the pair correlation function $g(r)$, and is only a function of Γ and the electron screening parameter $\kappa = \lambda_{\text{Debye}}/a$:

$$\frac{U_c}{NkT} \equiv \bar{U} = \frac{1}{2}\frac{e^2 n_0}{4\pi\epsilon_0}\int \frac{g(r,\Gamma,\kappa)d^3r}{r} \tag{2}$$

This binding energy is balanced by an increasing ion temperature. Calculating the increase in temperature thus only requires knowledge of $\bar{U}$ for a given electron screening, which have been tabulated via molecular dynamics (MD) simulations [9].

It has been demonstrated that a trapped, charged plasma, such as an electron bunch, is equivalent to the one-component plasma (OCP) model, in which the charges exist in a uniform neutralizing background [10]. The external containing potential allows the initial uncorrelated state to be viewed as zero total energy. As electrons relax, correlations develop, and the presence of a "Coulomb hole" in $g(r)$ for $r < a$ creates an effective correlation binding energy equal to that of a one component plasma. In the case of an OCP, there is no second species to provide additional screening, and thus $\bar{U}$ is only a function of Γ. Owing to this simplification, MD data has been fit to a power series relation for $\bar{U} = a\Gamma + b\Gamma^{1/3} + c$, where the coefficients are given in [10], $a = -0.90$, $b = 0.590673$, $c = -0.26569$. We may now write down the expression for the the final coupling Γ_f, and thus the final equilibrium temperature, for a fully confined, initially uncorrelated ($\bar{U}_i = 0$), uniform distribution of charges with initial coupling Γ_i, analogous to that presented in [9]:

$$\bar{U}_f = \frac{3}{2}\left(\frac{\Gamma_f}{\Gamma_i} - 1\right) = a\Gamma_f + b\Gamma_f^{1/3} + c \tag{3}$$

It is important to note the power series relation is quoted only for $\Gamma > 1$, however we find that for $\Gamma < 1$, the above expression is well approximated by $\Gamma_i = \Gamma_f$, or the limit of no DIH, as we expect. Eq. 3 is plotted in Fig. 1. For an ideal bunch with zero initial temperature, $\Gamma_i \to \infty$, Eq. 3 can be solved to give $\Gamma_f \approx 2.23$.

Practical relativistic photoelectron sources contain a number of complicating factors. Inside the emitting material (either a cold gas or crystal), the electron coupling, and thus DIH, will be dramatically reduced by the presence of the ionic/nuclear potential, and thus here we only need to consider DIH developing in vacuum. The beam density during emission will vary in time and with position across the bunch. The temperature associated with DIH should be reached on the order of ω_p^{-1} after electron emission into vacuum, which for typical beam densities is on the picosecond scale. On this scale, we can neglect relativistic effects, such as the $1/\gamma^2$ damping of the interaction force [5]. The coupling given by $\Gamma_f = 2.23 \equiv$

Γ_{eq}, for a given local beam density, can then be viewed as the fundamental photoemission temperature limit. We consider the effects of a changing local beam density, as well as the nonrelativistic effects of acceleration during the evolution of DIH below, and we find it to correspond well with the prediction of Eq. 3. Rewriting it in terms of practical units, the heating induced in a zero temperature bunch is given by $kT[\mathrm{eV}] = 1.04 \times 10^{-9}(n_0[\mathrm{m}^{-3}])^{1/3}$, which is only a function of interparticle separation. Along with Eq. 1, this forms the fundamental phoemission brightness limit.

To test the prediction of Eq. 3, we have chosen to use a tree-algorithm electrostatic simulation package, with nonrelativistic, spherical, uniform particle bunches. Traditional particle-in-cell integrators have an intrinsic length scale—the spatial grid on which local fields are calculated. However, tree algorithms lack a spatial grid, and thus are better suited to our initial conditions, where both close interactions and long range correlations are significant. Several plasma treecodes have been developed, see for instance [11]. However, considering the simplicity of the problem, we elected to directly modify a code originally written for gravitational interaction [12].

We consider a randomly distributed spherical bunch of 10^5 particles with uniform density, and vary only the density ($10^{17} \to 10^{20}$ m^{-3}) and initial Gaussian velocity distribution ($kT \in \{0.25, 20\}$ meV or $kT = 0$) , where we use the definition $kT = m_e \left(\langle v_i^2\rangle - \langle x_i v_i\rangle^2/\langle x_i^2\rangle\right)$, where m_e is the electron mass, and i is a Cartesian coordinate. There is no breaking of spherical symmetry in our simulations, thus equipartition will always hold. For a fully contained uniform bunch, $\langle x_i v_i\rangle \approx 0$, and the temperature is just a measure of average electron kinetic energy, with no spatial dependence. Containment is done in simulation via the application of a radial external force equal and opposite to the SC field. The parameters of the interparticle force calculation are the Barnes-Hut opening angle θ, and the leap frog integrator time step dt. The opening angle is the force calculation accuracy parameter that determines whether an electron is treated individually or lumped together with other similarly distant particles [13]. No effect was seen from a force-softening distance p, such that the force between two particles is $f_{12} \propto (r_{12} + p)^{-2}$, from $p = p = a/10^6$ up to $p = a/100$. For all simulations presented, we have demonstrated convergence using $dt = \tau/120$, $\theta = 23°$, $N = 10^5$.

For an initial distribution with zero temperature, the final temperature for several applicable beam densities is plotted in Fig. 2. We note that for extremely dense beams, $n_0 > 10^{19}$ m^{-3}, the DIH is comparable to the temperature of photoemission from the



FIG. 1: Final coupling Γ_f, vs initial coupling Γ_i, given by Eq. 3 (blue line), with $\Gamma_f = \Gamma_i$ for $\Gamma_i \to 0$ (red, dotted), and $\Gamma_f = 2.23$ for $\Gamma_i \to \infty$. Dots are simulation results of a fully contained, equipartitioned electron sphere, density $n_0 = 10^{20}$ m^{-3}, and initial temperatures between kT=0.25 meV and 20meV. Squares are simulation results for multiple densities, where $kT_z = 0$, and $kT_x = kT_y$, and where Γ is calculated from the average of the three directions. Error bars are an estimate of the uncertainty in final temperature determination due to the residual oscillations, as in Fig. 2.

coldest semiconductor cathodes $kT \sim 5$ meV. Furthermore, we note the oscillation of the electron temperature at $2\omega_p$, which has been measured experimentally in UNP systems [8]. These oscillations last far longer than those seen in UNPs, as the bunch remains uniform in simulation (rather than Gaussian), and there is thus no spatial temperature smearing.

For bunches with nonzero initial temperature, the amount of additional heating can be calculated in the final coupling, plotted along with the prediction of Eq. 3 in Fig. 1. All simulations here had a constant density $n_0 = 10^{20}$ m^{-3} , with varied initial Maxwellian velocity distribution. The duration of the temperature oscillations visible in Fig. 2 is reduced with increasing initial temperature, though the exact dependence was not extracted



FIG. 2: Temperature vs. number of plasma periods from treecode simulation using 10^5 contained electrons, and $n_0 = 10^{20}$ (top curve), 10^{19}, 10^{18}, 10^{17} m^{-3} (bottom curve), with zero initial temperature, $\Gamma_i \to \infty$. The equilibrium value predicted by Eq. 3, $\Gamma_f \approx 2.23$ is shown (dotted lines).

in this study. In practical systems, initial acceleration in a voltage gap V will have cooled any photocathode emission temperature along the acceleration direction by a factor kT_i/eV, making $kT_{||} \sim 0$ almost instantaneously in high gradient electron sources, whereas the transverse velocity spread will remain unaffected [5]. Simulations were performed for contained bunches of multiple densities with $kT_x = kT_y \neq 0$ and $kT_z = 0$. Equivalent dynamics to Fig. 2, as well as agreement with Eq. 3 were found if the average of the temperatures in all three directions is used to calculate Γ_i and Γ_f, as shown in Fig 1. The timescale of equipartition here also depends on kT_i and n_0, but for the densities considered above and initial temperatures comparable to DIH heating, it was seen that equipartition occurs between τ and 3τ. A bunch instantaneously emitted into a uniform accelerating field will not have the

DIH evolution altered, a fact that was also verified in simulation.

Real electron bunches just after emission are very infrequently fully transversely contained in the sense described above, and are not emitted instantaneously. The overall space charge force can double the beam radius on the timescale of ω_p^-1, and acceleration can significantly lengthen the pulse afterwards. Thus, we must consider the process of DIH in bunches with time dependent density reduction. A system of fully coupled charges will have an equilibrium temperature that decreases as $kT \propto 1/\Gamma_{eq}a$, as given by above. However, it is well known that system of noninteracting charges expanding under linear forces will have an equilibrium temperature that decreases as $kT \propto 1/a^2$ [5], as a consequence of RMS emittance conservation, which therefore leaves the phase space density or brightness unchanged. During the explosion, the beam is in a highly nonequilibrium state, but our definition of temperature above still applies, where here $\langle rv_r \rangle \neq 0$.

Simulations were performed to determine the temperature as a function of time for a bunch undergoing full Coulomb explosion. Multiple densities were considered, however the previous scaling with $n_0^{1/3}$ was seen. Thus, we present only results for $n_0 = 10^{20}$ m^{-3} in Fig. 3. The total overall expansion and velocity growth is plotted in the inset, with the analytic prediction, showing excellent agreement. The temperature shows an initial increase due to DIH at a time of $t_{\max} \approx \omega_p^{-1}$, to a value of $kT(t = t_{\max}) = e^2/4\pi\epsilon_0 a(t_{\max})$, or a value $a(t_{\max})/a_0 \approx 1.6$ times smaller than for a fully contained beam. The bunch continues to expand, attempting to equilibrate to $\Gamma_{eq} = 2.23$, and the temperature continues to decrease. However, at longer times, the temperature falls as a^{-2}, as in an uncoupled system. Thus, there must be a time at which correlation ceases to grow with decreasing temperature. Such behavior has been noted in the adiabatic expansion of UNPs due to kinetic pressure [14].

Thus, to model the temperature as a function of time, we can write the temperature as a product of the correlated growth, and the decoupled expansion after correlation ceases. Since the growth of the radius is small during the time of DIH, we may presume $T = T_c(t)\frac{a_0^2}{a(t)^2}$, where T_c is the correlated temperature dependence, and the term $\frac{a_0^2}{a(t)^2}$ describes the expansion of independent particles. The rate of DIH should be proportional to the how far the system is from the equilibrium temperature at the current density, and so:

$$\frac{dT_c}{dt} = \frac{T_{c,eq} - T_c(t)}{\tau_0} = \frac{1}{\tau_0}\left(\frac{e^2}{4\pi\epsilon_0 a(t)\Gamma_{eq}} - T_c(t)\right) \tag{4}$$

The only free parameter of this model is decoupling time τ_0, or the time when the correlation ceases to increase. Integrating this numerically, and fitting to the temperature data, we find $\tau_0 = 0.325\tau$, which is also plotted in Fig. 3. This time is non-negligibly greater than $t_{\max}$, and suggests that the bunch is cooled not only via decoupled expansion, but coupled expansion as well just after $t_{\max}$. We can verify this prediction by looking at the pair correlation as a function of time. This is computed in Fig. 4, via averaging over 5×10^3 particles randomly selected from the distribution. The development of the "Coulomb hole" and an accompanying shock profile due to violent repulsion is clearly visible, as has also been seen in UNP simulations [15]. Indeed the correlation ceases to change after a time of approximately $\tau_0 = 0.3\tau$. Partial confinement effected by linear focusing in the source would yield an altered $a(t)$, and thus a longer τ_0, whereas acceleration that lengthens the bunch significantly near the DIH heating timescale would yield a shorter τ_0, but in either case the DIH heating timescale should remain unchanged.

The exploding beam case justifies *post-factum* the validity of the temperature limit of Eq. 3 for practical photoemission, though beams in many applications do not have constant volume. In the exploding case, near τ_0 the temperature scaling shifts from $a(t)^{-1}$ as given in Eq. 2, to the familiar scaling of $\sigma_v^2 \propto a(t)^{-2}$, as given by emittance/brightness conservation. A more detailed calculation of DIH for practical cases in which the bunch shape is both time and space dependent should involve averaging over a temperature distribution given by Eqs. 3 and 4 using the local density as determined by space charge tracking.

In summary, this Letter has proposed that the fundamental temperature limit of photoemission will be from the disorder induced heating of electrons due to poorly screened Coulomb interactions at all length scales. We have shown that the tabulated thermodynamic quantities of one component plasmas are sufficient to explain both fully contained and Coulomb exploding instances of DIH, and have verified two simple relations that describe the temperature evolution. Furthermore, many interesting effects of DIH in UNP, such as temperature oscillation and correlation decoupling, should also be present in such cold beams, yielding the possibility of rich interdisciplinary study.

Practically, for next-generation ultracold dense electron sources we have shown that the temperature of photoemission, and thus the maximum beam brightness, cannot be arbitrarily improved. Furthermore, given the rapid progress of photocathode temperature reduction, we anticipate such a limit to be approached in the next generation of high brightness electron sources producing intense beams.



FIG. 3: RMS velocity spread of a spherical bunch undergoing coulomb expansion, treecode data (circles), and prediction of Eq. 4. The inset shows the overall expansion of the bunch in normalized units, both analytic prediction (solid line), given by Gauss's Law, and treecode data (dots). Uniform bunches remain uniform, and the normalized bunch size $R(\tau)/R_0$ is independent of density.

---

* Electronic address: jmm586@cornell.edu

[1] S. Pastuszka, D. Kratzmann, D. Schwalm, A. Wolf, and A. S. Terekhov, Applied Physics Letters **71**, 2967 (1997), URL `http://link.aip.org/link/?APL/71/2967/1`.

[2] Z. Liu, Y. Sun, P. Pianetta, and R. F. W. Pease (AVS, 2005), vol. 23, pp. 2758–2762, URL `http://link.aip.org/link/?JVB/23/2758/1`.

[3] G. Taban, M. P. Reijnders, B. Fleskens, S. B. van der Geer, O. J. Luiten, and E. J. D. Vredenbregt, EPL (Europhysics Letters) **91**, 46004 (2010), URL `http://stacks.iop.org/`



FIG. 4: Pair correlation function g(r), computed at multiple simulation times for a bunch undergoing coulomb explosion. Both the initial "shock" profile and the freezing out of correlations despite increasing Γ is visible.

`0295-5075/91/i=4/a=46004`.

[4] I. V. Bazarov, B. M. Dunham, and C. K. Sinclair, Phys. Rev. Lett. **102**, 104801 (2009), URL `http://link.aps.org/doi/10.1103/PhysRevLett.102.104801`.

[5] M. Reiser, *Theory and Design of Charged Particle Beams*, Wiley Series in Beam Physics and Accelerator Technology (John Wiley & Sons, 2008), ISBN 9783527622054, URL `http://books.google.com/books?id=eegK9Mqgpi4C`.

[6] G. H. Jansen, *Coulomb Interactions in Particle Beams*, Advances in Electronics & Electron Physics Supplement (Academic Press, 1990), ISBN 0120145839.

[7] M. Lyon and S. D. Bergeson, Journal of Physics B: Atomic, Molecular and Optical Physics **44**, 184014 (2011), URL `http://stacks.iop.org/0953-4075/44/i=18/a=184014`.

[8] Y. C. Chen, C. E. Simien, S. Laha, P. Gupta, Y. N. Martinez, P. G. Mickelson, S. B. Nagel,

and T. C. Killian, Phys. Rev. Lett. **93**, 265003 (2004), URL `http://link.aps.org/doi/10.1103/PhysRevLett.93.265003`.

[9] M. S. Murillo, Phys. Rev. Lett. **87**, 115003 (2001), URL `http://link.aps.org/doi/10.1103/PhysRevLett.87.115003`.

[10] D. H. E. Dubin and T. M. O'Neil, Rev. Mod. Phys. **71**, 87 (1999), URL `http://link.aps.org/doi/10.1103/RevModPhys.71.87`.

[11] B. Jeon, J. D. Kress, L. A. Collins, and N. Grnbech-Jensen, Computer Physics Communications **178**, 272 (2008), ISSN 0010-4655, URL `http://www.sciencedirect.com/science/article/pii/S0010465507004171`.

[12] J. Barnes, *Treecode guide* (2001), URL `http://www.ifa.hawaii.edu/~barnes/treecode/treeguide.html`.

[13] J. Barnes and P. Hut, Nature **324**, 446 (1986), URL `http://dx.doi.org/10.1038/324446a0`.

[14] T. Pohl, T. Pattard, and J. M. Rost, Phys. Rev. Lett. **94**, 205003 (2005), URL `http://link.aps.org/doi/10.1103/PhysRevLett.94.205003`.

[15] M. S. Murillo, Phys. Rev. Lett. **96**, 165001 (2006), URL `http://link.aps.org/doi/10.1103/PhysRevLett.96.165001`.